arXiv:1512.03297v1 [math.NT] 10 Dec 2015

STEINITZ CLASSES AND PARTIAL GENERA OF UNIMODULAR LATTICES OVER IMAGINARY-QUADRATIC FIELDS

MICHAEL JÜRGENS AND MARC C. ZIMMERMANN

Abstract. In this paper we first of all determine all possible genera of (odd and even) definite unimodular lattices over an imaginary-quadratic field. The main questions are whether the partial class numbers of lattices with given Steinitz class within one genus are equal for all occuring Steinitz classes and whether the partial masses of those partial genera are equal. We show that the answer to the first question in general is "no" by giving a counter example, while the answer to the second question is "yes" by proving a mass formula for partial masses. Finally, we determine a list of all single-class partial genera and show that a partial genus consists of only one class if and only if the whole genus consists of only one class.

Introduction

Let K be an algebraic number field. If L is a quadratic lattice over K, it is easily seen that the Steinitz class $\mathfrak{st}(L)$ is an invariant of the genus of L. In particular if one lattice in a genus is free so are all others. In general, this is not the case for hermitian lattices over K as the following three examples of genera of (positive) definite unimodular hermitian lattices over imaginary-quadratic fields show:

Example 1. Let $K := \mathbb{Q}(\sqrt{-47})$. The class group of K is the cyclic group of order 5 with generator $[\mathfrak{a}]$. There is exactly one genus of definite unimodular lattices of rank 4 over K:

		$[\mathfrak{o}]$	$[\mathfrak{a}]$	$[\mathfrak{a}^2]$	$[\mathfrak{a}^3]$	$[\mathfrak{a}^4]$	parity
		partial					
class number	1510	302	302	302	302	302	odd
mass	$\frac{1105}{8}$	$\frac{221}{8}$	$\frac{221}{8}$	$\frac{221}{8}$	$\frac{221}{8}$	$\frac{221}{8}$	

Example 2. Let $K := \mathbb{Q}(\sqrt{-95})$. The class group of K is the cyclic group of order 8 with generator $[\mathfrak{a}]$. In this case there are exactly two genera of definite unimodular lattices of rank 3 over K:

		$[\mathfrak{o}]$	$[\mathfrak{a}]$	$[\mathfrak{a}^2]$	$[\mathfrak{a}^3]$	$[\mathfrak{a}^4]$	$[\mathfrak{a}^5]$	$[\mathfrak{a}^6]$	$[\mathfrak{a}^7]$	parity
		partial								
class number	428	107	-	107	-	107	-	107	-	odd
mass	70	$\frac{35}{2}$	-	$\frac{35}{2}$	-	$\frac{35}{2}$	-	$\frac{35}{2}$	-	
class number	428	-	107	-	107	-	107	-	107	odd
mass	70	-	$\frac{35}{2}$	-	$\frac{35}{2}$	-	$\frac{35}{2}$	-	$\frac{35}{2}$	

Date: 11th December 2015.
2010 *Mathematics Subject Classification.* Primary <11E39,11E41>;Secondary <11R11>.
Key words and phrases. hermitian unimodular lattice, imaginary-quadratic field, mass formula, Steinitz class.

Example 3. Let $K := \mathbb{Q}(\sqrt{-213})$. The class group of K is the group $\mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ of order 8 with generators $[\mathfrak{a}]$ and $[\mathfrak{b}]$. There are exactly 4 genera of odd definite unimodular lattices and exactly 2 genera of even definite unimodular lattices of rank 2 over K:

		$[\mathfrak{o}]$	$[\mathfrak{a}]$	$[\mathfrak{b}]$	$[\mathfrak{ab}]$	$[\mathfrak{a}^2]$	$[\mathfrak{a}^3]$	$[\mathfrak{a}^2\mathfrak{b}]$	$[\mathfrak{a}^3\mathfrak{b}]$	parity
		partial								
class number	50	25	-	-	-	25	-	-	-	odd
mass	$\frac{35}{2}$	$\frac{35}{4}$	-	-	-	$\frac{35}{4}$	-	-	-	
class number	40	-	20	-	-	-	20	-	-	odd
mass	18	-	9	-	-	-	9	-	-	
class number	74	-	-	37	-	-	-	37	-	odd
mass	35	-	-	$\frac{35}{2}$	-	-	-	$\frac{35}{2}$	-	
class number	84	-	-	-	42	-	-	-	42	odd
mass	36	-	-	-	18	-	-	-	18	
class number	32	-	16	-	-	-	16	-	-	even
mass	12	-	6	-	-	-	6	-	-	
class number	56	-	-	28	-	-	-	28	-	even
mass	$\frac{70}{3}$	-	-	$\frac{35}{3}$	-	-	-	$\frac{35}{3}$	-	

These three examples as well as the explicit enumeration of some other genera of definite unimodular lattices of "low rank" over imaginary-quadratic fields with rather "small discriminant" lead to the following questions:

1. How many genera of definite unimodular lattices of given rank over a given imaginary-quadratic field do exist? In particular, which genera of even definite unimodular lattices do exist?
2. How do the Steinitz classes distribute among those genera? Does this distribution induce a partition of the class group?
3. Are the *partial class numbers* of lattices within a genus of definite unimodular lattices with the same Steinitz class equal for all occuring Steinitz classes?
4. Are the *partial masses* of lattices within a genus of definite unimodular lattices with the same Steinitz class equal for all occuring Steinitz classes?
5. What are the single-class partial genera of definite unimodular lattices?

In the first section of this paper we collect some basic definitions and facts about hermitian lattices over a number field K. We explain the notions of ideal genera as well as special classes and special genera of hermitian lattices. The latter lead to questions similar to 3 and 4 concerning special masses and special class numbers, which have been (partially) answered in the literature (c.f. [Reh71],[HK89a],[HK89b]). We describe a connection between special and partial genera, which we use in the following sections to deduce results about partial class numbers and partial masses from these known results.

In section 2 we cite a result of Hoffmann (c.f. [Hof91]) and Hashimoto and Koseki (c.f. [HK89a]) concerning the number of hermitian spaces carrying a definite unimodular lattice. Furthermore, we give a criterion for the existence of an odd resp. even definite unimodular lattice of given rank and given Steinitz class over a given imaginary-quadratic field. We deduce the essentially known result

that two definite unimodular lattices of same rank and parity are in the same genus if and only if their Steinitz classes are in the same ideal genus.
In section 3 we show that the answer to question 3 in general is "no" by giving a counter example. Actually, these counter examples seem to be very rare, at least for low rank and small discriminant. Therefore we try to find sufficient conditions on the rank and the class group that imply the equality of the partial class numbers in as many cases as possible. For example we can show that such an equality holds if the rank is 2 or the class group is 2-elementary. Using results of Hashimoto and Koseki (c.f. [HK89b]) we deduce the same result for lattices of rank 3.
In section 4 we use results of Rehmann (c.f. [Reh71]) to deduce a mass formula for partial masses which shows that the partial masses are indeed equal for all occuring Steinitz classes within a genus of definite unimodular lattices, i.e. the answer to question 4 is "yes". Furthermore, we evaluate the formulas in Gan and Yu (c.f. [GY00]) and Cho (c.f. [Cho12]) for the local densities of the genera determined in section 2 and we show how to compute these masses explicitly with the help of generalized Bernoulli numbers.
Finally, in section 5 we use the mass formula for the partial masses to compute a list of a single-class partial genera. It turns out that a partial genus of definite unimodular lattices consists of exactly one class if and only if the whole genus consists of only one class.

All computations described in this paper have been performed with Magma (c.f. [BCP97]). A description of the necessary algorithms, in particular the neighbour method for hermitian lattices can be found in [Sch98]. The implementations of these algorithms in Magma can be found under `http://www.mathematik.tu-dortmund.de/sites/michael-juergens/research`

1. Preliminaries

Throughout this paper let $K := \mathbb{Q}(\sqrt{D})$ be an imaginary-quadratic field, i.e. $D < 0$ and subsequently we will always assume that D is squarefree. We denote by $\overline{} : K \to K$, $\alpha \mapsto \overline{\alpha}$ the non-trivial galois automorphism of $K/\mathbb{Q}$. Furthermore, let $\mathfrak{o}$ be the ring of integers of K and d_K the discriminant of K. It is a well-known fact that $\mathfrak{o} = \mathbb{Z}[\frac{1+\sqrt{D}}{2}]$ and $d_K = D$ if $D \equiv_4 1$, and $\mathfrak{o} = \mathbb{Z}[\sqrt{D}]$ and $d_K = 4D$ if $D \equiv_4 2, 3$. Let $\mathbb{P}$ denote the set of rational prime numbers. A prime $p \in \mathbb{P}$ is called *ramified* (in the extension $K/\mathbb{Q}$) if there is a prime ideal $\mathfrak{p} \subseteq \mathfrak{o}$ with $p\mathfrak{o} = \mathfrak{p}^2$ and *unramified* otherwise. More precisely, an unramified prime p is called *inert* if $p\mathfrak{o} = \mathfrak{p}$ for some prime ideal $\mathfrak{p}$, and *split* if $p\mathfrak{o} = \mathfrak{p}\,\overline{mathfrakp}$ for some prime ideal $\mathfrak{p}$ with $\mathfrak{p} \neq \overline{\mathfrak{p}}$. Let χ_K denote the character associated to K that is the unique Dirichlet character satisfying

$$\chi_K(2) := \begin{cases} 0 & d_K \equiv_4 0 \\ 1 & d_K \equiv_8 1 \\ -1 & d_K \equiv_8 5 \end{cases} \quad \text{and} \quad \chi_K(p) := \left(\frac{d_K}{p}\right) \text{ for } p \neq 2.$$

It is also well-known that a prime $p \in \mathbb{P}$ is ramified (resp. inert/split) if and only if $\chi_k(p) = 0$ (resp. $-1/+1$). We denote by t the number of ramified primes in $K/\mathbb{Q}$, i.e. the number of distinct prime divisors of d_K. We set $\mathcal{C}_K$ to be the ideal class group of K, h_K its class number, and denote by $[\mathfrak{a}]$ the class of a (fractional) ideal $\mathfrak{a}$.

1.1. **Lattices and their genera.** By a *hermitian space* V over K we understand a K-vector space of dimension $n < \infty$ together with a hermitian form $h : V \times V \to K$ that is a sesquilinear form (linear in the first component) such that $h(x, y) = \overline{h(y, x)}$. Note that since $\overline{h(x, x)} = h(x, x)$, we have $h(x, x) \in \mathbb{Q}$ for all $x \in V$. A hermitian space V is called *(positive) definite*, if $h(x, x) > 0$ for all

$x \in V \setminus \{0\}$. We denote by $dV \in \mathbb{Q}^*/N_{\mathbb{Q}}^K(K^*)$ the *discriminant* of V, by $U(V)$ the *unitary group* of V, and we write $V \cong W$ for two *isometric* hermitian spaces. It is well-known that two definite hermitian spaces V and W are isometric if and only if $\dim(V) = \dim(W)$ and $dV = dW$ (c.f. [Sch85] Ch. 10, ex. 1.6).

A *(hermitian) lattice* L over K is a finitely generated $\mathfrak{o}$-module $L \subseteq V$ for some hermitian space V. We define the *rank* of L by $\mathrm{rk}(L) := \dim(K{\cdot}L)$ and call L a lattice *on* V if $K{\cdot}L = V$. L is called *definite* if $K{\cdot}L$ is. Every definite lattice L can be uniquely decomposed into indecomposable lattices $L_1, \ldots, L_s$ (but for their order):

$$L = L_1 \perp \ldots \perp L_s$$

(c.f. [Eic52] Satz 3). In particular we have the law of cancellation for definite lattices. On the other hand, every lattice L on V can be written as

$$L = \mathfrak{a}_1 x_1 + \ldots + \mathfrak{a}_n x_n, \tag{1}$$

with ideals $\mathfrak{a}_1, \ldots, \mathfrak{a}_n$ and a basis $x_1, \ldots, x_n$ of V. Let $\mathfrak{st}(L)$ denote the *Steinitz class* of L that is the class $[\mathfrak{a}] \in \mathcal{C}_K$ of the ideal $\mathfrak{a} := \prod_{i=1}^n \mathfrak{a}_i$, which is a well-defined module-invariant of L. Two lattices L and N are *isometric*, $L \cong N$, if a $\sigma \in U(V)$ exists with $\sigma(L) = N$. The set $\mathrm{cls}(L) := \{N \subseteq V \text{ an } \mathfrak{o}\text{-lattice } \mid \exists \sigma \in U(V) : N = \sigma(L)\}$ of isometric lattices is called the *(isometry) class* of L, and the set of $\sigma \in U(V)$ with $\sigma(L) = L$ is called the *unitary group* of L and denoted $U(L)$. We denote by $L^\# := \{x \in V \mid h(x, L) \subseteq \mathfrak{o}\}$ the *dual lattice* of L.

We define the *scale* $\mathfrak{s}L$ and the *norm* $\mathfrak{n}L$ of L to be the $\mathfrak{o}$-ideals generated by $\{h(x,y) \mid x, y \in L\}$ and $\{h(x,x) \mid x \in L\}$ respectively. For a lattice as in (1) we define the volume of L by $\mathfrak{v}L := \prod_{i=1}^n \mathfrak{a}_i\overline{\mathfrak{a}_i} \cdot \det(x_1, \ldots, x_n)$, the latter being the determinant of a Gram matrix of h relative to $x_1, \ldots, x_n$. This is an $\mathfrak{o}$-ideal independent of the choice of the basis and the ideals in (1) and thus well defined.

We call a lattice L on V $\mathfrak{a}$-*modular* for an ideal $\mathfrak{a}$ if $\mathfrak{s}L = \mathfrak{a}$ and $\mathfrak{v}L = \mathfrak{a}^n$. An $\mathfrak{o}$-modular lattice is called *unimodular*. It is easy to show that L is $\mathfrak{a}$-modular if and only if $\mathfrak{a}L^\# = L$ and in particular unimodular if and only if $L^\# = L$.

A lattice L is called *even* if $\mathfrak{n}L \subseteq 2\mathfrak{o}$, and *odd* otherwise, which is referred to as the *parity* of L. It is clear that $\mathfrak{n}L = a\mathfrak{o}$ for an $a \in \mathbb{Z}$, and $\mathrm{Tr}_{\mathbb{Q}}^K(\mathfrak{s}L)\mathfrak{o} \subseteq \mathfrak{n}L \subseteq \mathfrak{s}L$. Since

$$\mathrm{Tr}_{\mathbb{Q}}^K(\mathfrak{o}) = \begin{cases} \mathbb{Z} & d_K \text{ is odd,} \\ 2\mathbb{Z} & d_K \text{ is even,} \end{cases} \tag{2}$$

we have $\mathfrak{n}L = \mathfrak{o}$ or $\mathfrak{n}L = 2\mathfrak{o}$ for a unimodular lattice L. Thus a unimodular lattice is even if and only if $\mathfrak{n}L = 2\mathfrak{o}$, and odd if and only if $\mathfrak{n}L = \mathfrak{o}$.

For $p \in \mathbb{P}$ let $\mathbb{Q}_p$ be the localization of $\mathbb{Q}$ at p and $\mathbb{Z}_p$ its ring of integers. Let $K_p := K \otimes_{\mathbb{Q}} \mathbb{Q}_p$ and $\mathfrak{o}_p := \mathfrak{o} \otimes_{\mathbb{Z}} \mathbb{Z}_p$ the ring of integers of K_p. If p is inert or ramified, then $K_p \cong K_{\mathfrak{p}} = \mathbb{Q}_p(\sqrt{D})$ is a field and $\mathfrak{o}_p \cong \mathfrak{o}_{\mathfrak{p}}$, whereas we have $K_p \cong \mathbb{Q}_p \times \mathbb{Q}_p$ and $\mathfrak{o}_p \cong \mathbb{Z}_p \times \mathbb{Z}_p$ if p is split. For $p \in \mathbb{P}$ we make use of the *local norm residue symbol* $(\,\cdot\,, K/\mathbb{Q})_p : \mathbb{Q}_p^* \to \{\pm 1\}$ with $(\alpha, K/\mathbb{Q})_p = +1$ if and only if $\alpha \in N_{\mathbb{Q}_p}^{K_p}(K_p^*)$.

Set $V_p := V \otimes_{\mathbb{Q}} \mathbb{Q}_p$ equipped with the p-adic prolongation of h, and $L_p := L \otimes_{\mathbb{Z}} \mathbb{Z}_p \subseteq V_p$. The *genus* of a lattice L on V is defined to be

$$\mathrm{gen}(L) := \{N \subseteq V \text{ an } \mathfrak{o}\text{-lattice } \mid \forall p \in \mathbb{P} : L_p \cong M_p\}.$$

If $\mathcal{G}$ is a genus of unimodular lattices, the *class number* $h_{\mathcal{G}}$ is the number of isometry classes of lattices in $\mathcal{G}$ and if $L_1, \ldots, L_h$ is a set of representatives of the classes contained in $\mathcal{G}$, the *mass* of $\mathcal{G}$ is defined

as

$$\operatorname{mass}(\mathcal{G}) := \sum_{i=1}^{h_{\mathcal{G}}} \frac{1}{|U(L_i)|}.$$

If $[\mathfrak{a}] \in \mathcal{C}_K$ is an occuring Steinitz class of some lattice in $\mathcal{G}$, we denote the *partial genus* of all lattices in $\mathcal{G}$ with Steinitz class $[\mathfrak{a}]$ by $\mathcal{G}_{[\mathfrak{a}]}$. For a partial genus $\mathcal{G}_{[\mathfrak{a}]}$ we denote its *partial class number* by $h_{\mathcal{G}_{[\mathfrak{a}]}}$. As a shorthand notation, we will sometimes use $h_{[\mathfrak{a}]}$ if the associated genus is clear from context. Assuming that $L_1, \dots, L_{h_{[\mathfrak{a}]}}$ is a set of representatives for $\mathcal{G}_{[\mathfrak{a}]}$, the *partial mass* of $\mathcal{G}_{[\mathfrak{a}]}$ is defined by

$$\operatorname{mass}(\mathcal{G}_{[\mathfrak{a}]}) := \sum_{i=1}^{h_{[\mathfrak{a}]}} \frac{1}{|U(L_i)|}.$$

1.2. **Relation between special and partial genera.** When replacing the unitary group by the special unitary group in the above definitions one derives the analogous notions of *special class*, *special genus*, *special class number* and *special mass*. They will be denoted by $\operatorname{cls}^+, \operatorname{gen}^+$, h^+ and mass^+.

It is easy to see that the special classes also put a partition on the set of all lattices on V, and that this partition is finer than the partition into isomety classes. Similarly, the partition into special genera is finer than the one into genra. We start by describing how a class partitions into special classes.

Lemma 1.1. *The class of L is the disjoint union of exactly $e(L) := [\mathfrak{o}^* : \det U(L)]$ special classes.*

Proof. The group $\mathfrak{o}^* \cong U(V)/SU(V)$ acts transitively on the special classes contained in the class of L, and an easy calculation shows that the stabilizer of this action is exactly $\det U(L) \cong \{\sigma SU(V) \mid \det \sigma \in \det U(L)\}$. □

We will now specify the relation between partial and special genera.

Lemma 1.2 (c.f. [HK89a], Lemma 2.4)**.** *A partial genus $\mathcal{G}_{[\mathfrak{a}]}$ is the disjoint union of exactly $e(\mathcal{G})$ special genera, where $e(\mathcal{G}) := \prod_{p \in \mathbb{P}} [\mathfrak{o}_p^{(1)} : \det U(L_p)]$ and $\mathfrak{o}_p^{(1)} := \{\varepsilon \in \mathfrak{o}_p^* \mid \varepsilon\overline{\varepsilon} = 1\}$. Furthermore, $e(\mathcal{G}) \leq 2$, with $e(\mathcal{G}) = 1$ if the rank or the parity is odd.* □

Corollary 1.3. *If $d_K \notin \{-3, -4\}$, special and partial class numbers coincide for lattices of odd rank.*

Proof. Let L be a lattice of odd rank n. We note that $-\operatorname{id} \in U(L)$ and $\det(-\operatorname{id}) = (-1)^n = -1$. Since $d_K \notin \{-3, -4\}$, we have $|\mathfrak{o}^*| = 2$, and therefore $[\mathfrak{o}^* : \det U(L)] = 1$. By Lemma 1.1 the class and the special class of L coincide. Now Lemma 1.2 gives $\mathcal{G}_{[\mathfrak{a}]} = \mathcal{G}^+$, which is exactly our claim. □

1.3. **Ideal genera and genus characters.** A *genus character* of K is a real-valued character of the class group that is a homomorphism $\chi : \mathcal{C}_K \to \{\pm 1\}$.

Proposition 1.4 (c.f. [Zag81] §12 Satz 1)**.** *Let $[\mathfrak{a}], [\mathfrak{b}] \in \mathcal{C}_K$, then the following three statements are equivalent:*

a) $\chi([\mathfrak{a}]) = \chi([\mathfrak{b}])$ for every genus character χ of K.
b) There is a $\lambda \in K$ with $N(\mathfrak{a}) = N(\lambda) \cdot N(\mathfrak{b})$.
c) There is a $[\mathfrak{c}] \in \mathcal{C}_K$ with $[\mathfrak{a}] = [\mathfrak{c}]^2 \cdot [\mathfrak{b}]$. □

If the equivalent conditions of 1.4 are fulfilled we say that $[\mathfrak{a}], [\mathfrak{b}] \in \mathcal{C}_K$ belong to the same *ideal genus*. The set of all ideal classes in the same genus as $[\mathfrak{a}]$ is denoted by $[\![[\mathfrak{a}]]\!]$ or simply by $[\![\mathfrak{a}]\!]$.

Since $|\mathcal{C}_K/\mathcal{C}_K^2| = 2^{t-1}$ (c.f. [Zag81] §12 Korollar), there are exactly 2^{t-1} ideal genera and since ideal genera are precisely the cosets of $\mathcal{C}_K^2$ in $\mathcal{C}_K$, all ideal genera contain the same number of classes,

namely $\frac{h_K}{2^{t-1}}$. Furthermore, the set of genus characters is precisely the dual group of the (abelian) group $\mathcal{C}_K/\mathcal{C}_K^2$ and therefore there are exactly 2^{t-1} genus characters.

Lemma 1.5. *Let $[\mathfrak{a}] \in \mathcal{C}_K$. Then we have $(N(\mathfrak{a}), K/\mathbb{Q})_p = 1$ for all unramified primes $p \in \mathbb{P}$ and moreover*

$$\prod_{p \text{ ramified}} (N(\mathfrak{a}), K/\mathbb{Q})_p = 1.$$

Proof. It is well known that in the case of a quadratic field $(\alpha, K/\mathbb{Q})_p = (\alpha, D)_p$, the latter being the Hilbert symbol at p. Let p be an unramified prime, thus D is a unit in $\mathbb{Q}_p$. Because there is always an integral ideal in $[\mathfrak{a}]$ which is coprime to $p\mathfrak{o}$ we can assume that $\gcd(N(\mathfrak{a}), p) = 1$. For $p \neq 2$ we are done, because the Hilbert symbol of two units always equals 1 in that case. If $p = 2$ we have $D \equiv_4 1$, which now implies the same as before.

The product formula now follows immediately from the Hilbert reciprocity law. □

Now consider the character $\chi_p : \mathcal{C}_K \to \{\pm 1\}$, $[\mathfrak{a}] \mapsto (N(\mathfrak{a}), K/\mathbb{Q})_p$, which is a genus character for every $p \in \mathbb{P}$. By the above Lemma we have $\chi_p \equiv 1$ for every unramified p.

The image of the homomorphism

$$\mathcal{X} : \mathcal{C}_K \to \{\pm 1\}^t,\ [\mathfrak{a}] \mapsto \left(\chi_p\left([\mathfrak{a}]\right)\right)_{p \text{ ramified}},$$

is a subset of $\mathcal{H} := \{(x_1, \ldots, x_t) \in \{\pm 1\}^t \mid \prod x_i = 1\}$. To determine the kernel of $\mathcal{X}$, let $[\mathfrak{a}] \in \ker \mathcal{X}$ be arbitrarily chosen, then $(N(\mathfrak{a}), K/\mathbb{Q})_p = 1$ for all p, so that $N(\mathfrak{a})$ is a norm for each $p \in \mathbb{P}$. By the Hasse norm Theorem (c.f. [O'M73] 65:23), there exists $\lambda \in K$ such that $N(\mathfrak{a}) = N(\lambda) = N(\lambda) \cdot N(\mathfrak{o})$. By Proposition 1.4 this is equivalent to $[\mathfrak{a}] = [\mathfrak{c}]^2$ for some $[\mathfrak{c}] \in \mathcal{C}_K$, thus $[\mathfrak{a}] \in \mathcal{C}_K^2$ and $\ker \mathcal{X} \subseteq \mathcal{C}_K^2$. The inclusion $\mathcal{C}_K^2 \subseteq \ker \mathcal{X}$ is obvious, therefore $\mathcal{X}$ induces an injective homomorphism:

$$\overline{\mathcal{X}} : \mathcal{C}_K/\mathcal{C}_K^2 \hookrightarrow \mathcal{H}.$$

Since there are 2^{t-1} ideal genera, i.e. $|\mathcal{C}_K/\mathcal{C}_K^2| = 2^{t-1}$, the map $\overline{\mathcal{X}}$ is bijective. In particular, the map $\mathcal{X} : \mathcal{C}_K \to \mathcal{H}$ is surjective and we have proved:

Proposition 1.6. *If d_K is even, $d_K \notin \{-4, -8\}$, there is an ideal class $[\mathfrak{a}] \in \mathcal{C}_K$, such that*

$$(N(\mathfrak{a}), K/\mathbb{Q})_2 = -1.$$

Proof. The condition that d_K is even and $d_K \notin \{-4, -8\}$ implies that 2 is ramified and $t \geq 2$. Since $\mathcal{X} : \mathcal{C}_K \to \mathcal{H}$ is surjective, there is a $[\mathfrak{a}] \in \mathcal{C}_K$ with $\mathcal{X}([\mathfrak{a}]) = (-1, -1, 1, \ldots, 1)$, i.e. $\chi_2([\mathfrak{a}]) = -1$. □

2. Genera of unimodular lattices

For the sake of self containment we start with the characterization those hermitian spaces that carry unimodular lattices, which is a well-known result. We also derive some necessary (and indeed sufficient as Theorem 2.3 will show) conditions for a hermitian space to carry an even unimodular lattice.

Proposition 2.1. *Let $K = \mathbb{Q}(\sqrt{D})$ and $n \in \mathbb{N}$.*

a) There are exactly 2^{t-1} definite hermitian spaces of dimension n over K (up to isometry) containing a unimodular lattice of rank n. There is a bijection between those hermitian spaces and the ideal genera in $\mathcal{C}_K$ given by $V \mapsto [\![\mathfrak{st}(L)]\!]$ where L is any unimodular lattice L on V.

b) For each definite hermitian space V there are at most two genera of unimodular lattices on V. For each parity there is at most one such genus.

c) *For the existence of an even definite unimodular lattice L of rank n on V, with $\mathfrak{st}(L) = [\mathfrak{a}]$, the following restrictions apply:*
   i) d_K must be even.
   ii) n must be even.
   iii) If $D \equiv_4 3$ then $(N(\mathfrak{a}), K/\mathbb{Q})_2 = (-1)^{\frac{n}{2}}$.

Proof. a) See [Hof91] 3.7 or [HK89a] 6.4. b) See [HK89a] 6.4. c) Formula (2) shows that d_K is necessarily even and therefore $p = 2$ is ramified. By [Jac62] Proposition 10.3 we have $L_2 \cong \mathbb{H} \perp \ldots \perp \mathbb{H} \perp N$, where N is a unimodular $\mathfrak{o}_2$-lattice of rank 1 or 2 with $\mathfrak{n}L_2 = \mathfrak{n}N$. Since there are no even unimodular $\mathfrak{o}_2$-lattices of rank 1, we conclude $\mathrm{rk}(N) = 2$. This implies that n is even. If $D \equiv_4 3$ we are in the ramified-unit (RU) case (c.f. [Jac62] Chapter 5), for which [Jac62] Proposition 9.2 shows that $N \cong \mathbb{H}$ that is $(-1)^{\frac{n}{2}} \in dV_2$. On the other hand $N(\mathfrak{a}) \in dV$, since $\mathfrak{o} = \mathfrak{v}L = \mathfrak{a}\overline{\mathfrak{a}} \cdot \det(x_1, x_2)$, and $\det(x_1, x_2), N(\mathfrak{a})$ are positive rational numbers, we conclude $N(\mathfrak{a}) \cdot \det(x_1, x_2) = 1$. Therefore

$$(N(\mathfrak{a}), K/\mathbb{Q})_2 = ((-1)^{\frac{n}{2}}, K/\mathbb{Q})_2 = ((-1)^{\frac{n}{2}}, D)_2 = (-1)^{\frac{n}{2}},$$

where $D \equiv_4 3$ implies that $(-1, D)_2 = (-1, 3)_2 = -1$ or $(-1, D)_2 = (-1, 7)_2 = -1$. □

The 2^{t-1} definite hermitian spaces containing a unimodular lattice are given by $\langle 1, \ldots, 1, \frac{1}{N(\mathfrak{a})} \rangle$, where $[\mathfrak{a}]$ runs through a set of representatives of the 2^{t-1} ideal genera.
As an easy consequence we get:

Corollary 2.2. *Let L, N be two definite unimodular lattices of the same rank and parity. Then L and N are in the same genus if and only if $\mathfrak{st}(L)$ and $\mathfrak{st}(N)$ are in the same ideal genus, i.e.*

$$\mathrm{gen}(L) = \mathrm{gen}(N) \quad \Leftrightarrow \quad [\![\mathfrak{st}(L)]\!] = [\![\mathfrak{st}(N)]\!].$$

Proof. If L and N are lattices in the same genus, they are lattices on the same hermitian space. By Prop. 2.1a) the Steinitz classes of L and N are in the same genus. On the other hand if L and N are unimodular lattices of the same rank with $[\![\mathfrak{st}(L)]\!] = [\![\mathfrak{st}(N)]\!]$, by Prop. 2.1a) L and N are lattices on the same hermitian space. Since the parity of L and N coincides, L and N are in the same genus by Prop. 2.1b). □

Thus a genus of definite unimodular lattices over an imaginary-quadratic field is entirely determined by the parity, rank and ideal genus of the Steinitz class of any lattice in it. We denote the genus corresponding to the parity odd resp. even, the rank $n \in \mathbb{N}$ and the ideal genus $[\![\mathfrak{a}]\!]$ by $\mathbb{I}_{n,[\![\mathfrak{a}]\!]}$ resp. $\mathbb{II}_{n,[\![\mathfrak{a}]\!]}$.

We will now show that the necessary conditions of Proposition 2.1 are also sufficient, i.e. we will give a construction for a sample lattice $L^{\mathrm{odd}}_{n,[\mathfrak{a}]}$ resp. $L^{\mathrm{even}}_{n,[\mathfrak{a}]}$ in each possible genus of (odd or even) definite unimodular lattices.

Theorem 2.3. *Let $[\mathfrak{a}] \in \mathcal{C}_K$ and $n \in \mathbb{N}$.*

a) *There is a definite odd unimodular lattice of rank n with Steinitz class $[\mathfrak{a}]$ over K.*
b) *There is a definite even unimodular lattice of rank n with Steinitz class $[\mathfrak{a}]$ over K if and only if the following conditions are satisfied:*
   i) d_K is even.
   ii) n is even.
   iii) If $D \equiv_4 3$, then $(N(\mathfrak{a}), K/\mathbb{Q})_2 = (-1)^{\frac{n}{2}}$

Proof. a) The lattice $L^{\mathrm{odd}}_{1,[\mathfrak{a}]} := \mathfrak{a}x_1$ on the hermitian space $\langle \frac{1}{N(\mathfrak{a})} \rangle$ (in the basis x_1) is clearly an odd definite unimodular lattice of rank 1 with Steinitz class $[\mathfrak{a}]$. Therefore the lattice

$$L^{\mathrm{odd}}_{n,[\mathfrak{a}]} := L^{\mathrm{odd}}_{1,[\mathfrak{o}]} \perp \ldots \perp L^{\mathrm{odd}}_{1,[\mathfrak{o}]} \perp L^{\mathrm{odd}}_{1,[\mathfrak{a}]}$$

is an odd definite unimodular lattice of rank n and Steinitz class $[\mathfrak{a}]$.

b) The necessity of the conditions is an immediate consequence of 2.1*c*). Let $[\mathfrak{a}]$ be an ideal class as allowed by the above restrictions. We start with $n = 2$:
W.l.o.g. assume that $\mathfrak{a}$ is an integral ideal such that $2 \nmid N(\mathfrak{a})$. We claim that there exist $\lambda \in \mathbb{N}$ and $\alpha \in \mathfrak{o}$, such that $L := \mathfrak{o}x_1 + \mathfrak{a}x_2$ on the hermitian space given by the Gram matrix $\begin{pmatrix} 2 & \alpha \\ \overline{\alpha} & \frac{2\lambda}{N(\mathfrak{a})} \end{pmatrix}$ (in the basis x_1, x_2) is a binary even unimodular lattice with Steinitz class $[\mathfrak{a}]$: For $N(\mathfrak{a}) \equiv_4 -1$ we choose $\alpha := 1$ and $\lambda := \frac{1+N(\mathfrak{a})}{4}$ and for $N(\mathfrak{a}) \equiv_4 1$ we choose $\alpha := 1 + \sqrt{D}$ and $\lambda := \frac{1+(1+D)N(\mathfrak{a})}{4}$. In the latter case conditions *i*) and *iii*) imply $D \equiv_4 2$, so that in either case $\lambda \in \mathbb{N}$. Now consider the volume and scale of L:

$$\mathfrak{v}L = \mathfrak{a}\overline{\mathfrak{a}} \cdot \det(x_1, x_2) = \mathfrak{a}\overline{\mathfrak{a}} \cdot \tfrac{1}{N(\mathfrak{a})} = \mathfrak{o}, \quad \mathfrak{s}L = 2\mathfrak{o} + \alpha\overline{\mathfrak{a}} + \overline{\alpha}\mathfrak{a} + \tfrac{2\lambda}{N(\mathfrak{a})}\mathfrak{a}\overline{\mathfrak{a}} \subseteq \mathfrak{o}.$$

Since $\mathfrak{o} = \mathfrak{v}L \subseteq (\mathfrak{s}L)^2 \subseteq \mathfrak{o}$, we conclude $\mathfrak{s}L = \mathfrak{o}$. Therefore L is a unimodular lattice. Since

$$\mathfrak{n}L = 2\mathfrak{o} + 2\lambda\mathfrak{o} + \mathrm{Tr}^K_{\mathbb{Q}}(\mathfrak{o}) = 2\mathfrak{o},$$

L is even. Clearly the Steinitz class of L is $[\mathfrak{a}]$ and L is definite. Set $L^{\mathrm{even}}_{2,[\mathfrak{a}]} := L$.

For arbitrary even n we split the proof depending on D mod 4. For $D \equiv_4 2$ we construct $L^{\mathrm{even}}_{2,[\mathfrak{a}]}$ as above. To obtain lattices of rank $n = 2m$ we construct

$$L^{\mathrm{even}}_{n,[\mathfrak{a}]} := L^{\mathrm{even}}_{2,[\mathfrak{o}]} \perp \ldots \perp L^{\mathrm{even}}_{2,[\mathfrak{o}]} \perp L^{\mathrm{even}}_{2,[\mathfrak{a}]}.$$

For $D \equiv_4 3$, we distinguish between the possible values of the norm residue symbol for $N(\mathfrak{a})$. If $(N(\mathfrak{a}), K/\mathbb{Q})_2 = -1$, then *iii*) implies $n = 4m + 2$. We construct

$$L^{\mathrm{even}}_{n,[\mathfrak{a}]} := (L^{\mathrm{even}}_{2,[\mathfrak{a}]} \perp L^{\mathrm{even}}_{2,[\mathfrak{a}^{-1}]}) \perp \ldots \perp (L^{\mathrm{even}}_{2,[\mathfrak{a}]} \perp L^{\mathrm{even}}_{2,[\mathfrak{a}^{-1}]}) \perp L^{\mathrm{even}}_{2,[\mathfrak{a}]}.$$

If $(N(\mathfrak{a}), K/\mathbb{Q})_2 = 1$, then *iii*) implies $n = 4m$. If $D \neq -1$, Corollary 1.6 implies that we can choose another ideal class $[\mathfrak{b}]$ with $(N(\mathfrak{b}), K/\mathbb{Q})_2 = -1$ and construct

$$L^{\mathrm{even}}_{n,[\mathfrak{a}]} := (L^{\mathrm{even}}_{2,[\mathfrak{b}]} \perp L^{\mathrm{even}}_{2,[\mathfrak{b}^{-1}]}) \perp \ldots \perp (L^{\mathrm{even}}_{2,[\mathfrak{b}]} \perp L^{\mathrm{even}}_{2,[\mathfrak{b}^{-1}]}) \perp L^{\mathrm{even}}_{2,[\mathfrak{b}]} \perp L^{\mathrm{even}}_{2,[\mathfrak{b}^{-1}\mathfrak{a}]}.$$

Finally if $D = -1$, let $L^{\mathrm{even}}_{4,[\mathfrak{o}]} := \mathfrak{o}x_1 + \mathfrak{o}x_2 + \mathfrak{o}x_3 + \mathfrak{o}x_4$ on the hermitian space given by the following Gram matrix (in the basis $x_1, \ldots, x_4$):

$$\begin{pmatrix} 2 & 1 & 0 & 0 \\ 1 & 2 & 1+i & 1+i \\ 0 & 1-i & 2 & 1 \\ 0 & 1-i & 1 & 2 \end{pmatrix}$$

It is easily verified that $L^{\mathrm{even}}_{4,[\mathfrak{o}]}$ is an even definite unimodular lattice. For arbitrary $n = 4m$ we construct

$$L^{\mathrm{even}}_{n,[\mathfrak{o}]} := L^{\mathrm{even}}_{4,[\mathfrak{o}]} \perp \ldots \perp L^{\mathrm{even}}_{4,[\mathfrak{o}]}.$$

In every case the lattice $L^{\mathrm{even}}_{n,[\mathfrak{a}]}$ is an even definite unimodular lattice with Steinitz class $[\mathfrak{a}]$ as claimed. □

Corollary 2.4. *Let K be an imaginary-quadratic field such that d_K is even. Then there is a free quaternary even definite unimodular lattice over K.* □

3. Distribution of Steinitz classes

In this section we investigate the question whether $h_{[\mathfrak{a}]} = h_{[\mathfrak{b}]}$ for $[\mathfrak{a}], [\mathfrak{b}] \in \mathcal{C}_K$ holds. Since ideals and (odd) lattices of rank 1 happen to be essentially the same objects, we have $h_{[\mathfrak{a}]} = 1$ for all $\mathfrak{a} \in \mathcal{C}_K$ and therefore the problem is trivial in this case. For higher rank, we can show that in many cases such an equality arises from algebraic constructions. In particular in the binary case such a relation holds for all $[\mathfrak{a}], [\mathfrak{b}]$ in the same ideal genus as Corollary 3.7 will show. Using results of [HK89b], we can deduce the same result in the case ternary lattices. One might expect that this phenomenon is true in general, but the following counter-example for a genus of rank 4 shows that this is not the case:

Example 3.1. Let $K = \mathbb{Q}(\sqrt{-39})$. The class group of K is the cyclic group of order 4 with generator $[\mathfrak{a}]$. Because $e(\mathcal{G}) = 1$ (c.f. Lemma 1.2) for both genera $\mathcal{G}$ of odd definite unimodular lattices of rank 4, the special genera coincide with the partial genera. In the following table, they are indexed by the Steinitz class of the lattices they contain:

		$\mathcal{G}_{[\mathfrak{o}]}$	$\mathcal{G}_{[\mathfrak{a}]}$	$\mathcal{G}_{[\mathfrak{a}^2]}$	$\mathcal{G}_{[\mathfrak{a}^3]}$	$\mathcal{G}_{[\mathfrak{o}]}$	$\mathcal{G}_{[\mathfrak{a}]}$	$\mathcal{G}_{[\mathfrak{a}^2]}$	$\mathcal{G}_{[\mathfrak{a}^3]}$
		partial				special			
$\mathbb{I}_{4,[\![\mathfrak{o}]\!]}$	class number	162	-	164	-	172	-	176	-
	mass	$\frac{935}{72}$	-	$\frac{935}{72}$	-	$\frac{935}{36}$	-	$\frac{935}{36}$	-
$\mathbb{I}_{4,[\![\mathfrak{a}]\!]}$	class number	-	152	-	152	-	152	-	152
	mass	-	$\frac{154}{15}$	-	$\frac{154}{15}$	-	$\frac{308}{15}$	-	$\frac{308}{15}$

The above example is a minimal one in the sense that the discriminant $d_K = -39$ is the discriminant of smallest absolute value among all discriminants of imaginary-quadratic fields for which there is a genus of definite unimodular lattices containing partial genera with non-equal partial class numbers. There are also examples of even unimodular lattices of rank 4 where such a phenomenon occurs, like e.g. the genus $\mathbb{III}_{4,[\![\mathfrak{o}]\!]}$.

We first of all want to explain the equality of the partial class numbers within the genus $\mathbb{I}_{4,[\![\mathfrak{a}]\!]}$. We denote by $\overline{L}$ the lattice $\overline{\mathfrak{a}}_1 x_1 + \ldots + \overline{\mathfrak{a}}_n x_n$ on the hermitian space $(V, \overline{h})$, where $\overline{h}(x,y) := \overline{h(x,y)}$.

Lemma 3.2. *The map $L \mapsto \overline{L}$ maps isometry classes of definite unimodular lattices to isometry classes of definite unimodular lattices preserving the rank, the parity and the unitary group.* □

Proposition 3.3. *Let $\mathcal{G}$ be a genus of definite unimodular lattices of rank n over K and $[\mathfrak{a}] \in \mathcal{C}_K$. Then the map $L \mapsto \overline{L}$ induces a bijection between $\mathcal{G}_{[\mathfrak{a}]}$ and $\mathcal{G}_{[\overline{\mathfrak{a}}]}$, i.e. $h_{[\mathfrak{a}]} = h_{[\overline{\mathfrak{a}}]}$.* □

Now, let $L = \mathfrak{a}_1 x_1 + \ldots + \mathfrak{a}_n x_n$ be a unimodular lattice on the hermitian space (V, h). Denote by ${}^{\mathfrak{a}}L$ the lattice $\mathfrak{a}L$ on the hermitian space $(V, \frac{1}{N(\mathfrak{a})}h)$. The following lemma is well-known and straightforward to check (c.f. [Hof91]):

Lemma 3.4. *The map $L \mapsto {}^{\mathfrak{c}}L$ maps isometry classes of definite unimodular lattices to isometry classes of definite unimodular lattices preserving the rank, the parity and the unitary group.* □

Proposition 3.5. *Let $\mathcal{G}$ and $\mathcal{G}'$ be two genera of definite unimodular lattices over K of rank n and same parity. If there is a $[\mathfrak{c}] \in \mathcal{C}_K$ with $[\mathfrak{b}] = [\mathfrak{c}]^n \cdot [\mathfrak{a}]$, then the map $L \mapsto {}^{\mathfrak{c}}L$ induces a bijection between $\mathcal{G}_{[\mathfrak{a}]}$ and $\mathcal{G}'_{[\mathfrak{b}]}$, i.e. $h_{[\mathfrak{a}]} = h_{[\mathfrak{b}]}$.*

Proof. One immediately sees that $\mathfrak{st}({}^{\mathfrak{c}}L) = [\mathfrak{c}]^n \mathfrak{st}(L)$ where n is the rank of L. The rest is covered by the above Lemma. □

Theorem 3.6. *For definite unimodular lattices of fixed rank n and fixed parity we have:*

a) If $\gcd(n, h_K) = 1$*, then* $h_{[\mathfrak{a}]} = h_{[\mathfrak{b}]}$ *for all* $[\mathfrak{a}], [\mathfrak{b}] \in \mathcal{C}_K$*.*

b) If $\gcd(n, \exp(\mathcal{C}_K)) \leq 2$*, then* $h_{[\mathfrak{a}]} = h_{[\mathfrak{b}]}$ *for all* $[\mathfrak{b}] \in [\![\mathfrak{a}]\!]$*.*

Proof. *a*) Using Lemma 3.5, we have to show that for all $[\mathfrak{a}], [\mathfrak{b}] \in \mathcal{C}_K$ there exists $[\mathfrak{c}] \in \mathcal{C}_K$, such that $[\mathfrak{b}] = [\mathfrak{c}]^n \cdot [\mathfrak{a}]$, which is equivalent to $\mathcal{C}_K^n = \mathcal{C}_K$. Now the assumption $\gcd(n, h_K) = 1$ implies this equality, as taking n-th powers becomes an automorphism of the finite abelian group $\mathcal{C}_K$.

b) We have to show that for all $[\mathfrak{b}] \in [\![\mathfrak{a}]\!]$ there is a $[\mathfrak{c}] \in \mathcal{C}_K$ with $[\mathfrak{b}] = [\mathfrak{c}]^n \cdot [\mathfrak{a}]$. Since $[\mathfrak{a}]$ and $[\mathfrak{b}]$ are in the same ideal genus, this is equivalent to $\mathcal{C}_K^2 \subseteq \mathcal{C}_K^n$, which is essentially a group theoretic question.

We use a decomposition of $\mathcal{C}_K$ into cyclic p-groups. Since $\exp(\mathcal{C}_K)$ is precisely the product of the largest orders of those cyclic factors, it follows from the assumption $\gcd(n, \exp(\mathcal{C}_K)) \leq 2$ that $\gcd(n, |C|) \leq 2$ for each such C. For p odd or $p = 2$ and n odd we have $C^n = C$ by the same argument as in a) and therefore $C^2 \subseteq C = C^n$ in these cases. Now assume that $p = 2$ and n is even. Then $|C| \leq 2$ or $\frac{n}{2}$ is odd. But then $C^2 = \{1\} \subseteq C^n$ is trivially in the first case and in the second case we have $\gcd(\frac{n}{2}, |C|) = 1$, which then implies $C^{\frac{n}{2}} = C$, thus $C^n = C^2$.

Now since $C^2 \subseteq C^n$ for every cyclic factor, we conclude $\mathcal{C}_K^2 \subseteq \mathcal{C}_K^n$ as proposed. □

We now consider the two trivial cases in which the condition of part *b*) of the above Theorem is satisfied.

Corollary 3.7. *The isometry classes of definite unimodular lattices of rank n in the same genus, with given Steinitz classes $[\mathfrak{a}], [\mathfrak{b}]$, are in bijection if*

i) $\mathcal{C}_K$ is 2-elementary, or

ii) $n = 2$.

Examples not covered by Theorem 3.6 respectively Corollary 3.7 are the following:

Example 3.8. a) Let $K = \mathbb{Q}(\sqrt{-23})$. The class group of K is the cyclic group of order 3 with generator $[\mathfrak{a}]$. The genus $\mathbb{I}_{3,[\mathfrak{o}]}$ is partitioned into 3 times 10 isometry classes with respectively equal Steinitz classes. However, if one counts the number of classes in the three partial genera with same decomposition type, the following table is derived:

	1+1+1	2+1	3
$\mathcal{G}_{[\mathfrak{o}]}$	4	3	3
$\mathcal{G}_{[\mathfrak{a}]}$	3	3	4
$\mathcal{G}_{[\mathfrak{a}]^2}$	3	3	4

b) Let $K = \mathbb{Q}(\sqrt{-83})$, so $\mathfrak{o} = \mathbb{Z}[\frac{1+\sqrt{-83}}{2}]$. The class group of K is the cyclic group of order 3 with generator $[\mathfrak{a}]$. The genus $\mathbb{I}_{3,[\mathfrak{o}]}$ is partitioned into 3 times 43 isometry classes with respectively equal Steinitz classes, however the sequence of orders of the associated unitary groups do only match for two of these Steinitz classes:

	2	4	8	12	16	24	48
$\mathcal{G}_{[\mathfrak{o}]}$	6	21	10	0	0	3	3
$\mathcal{G}_{[\mathfrak{a}]}$	7	18	9	3	3	3	0
$\mathcal{G}_{[\mathfrak{a}]^2}$	7	18	9	3	3	3	0

In contrast to the bijections $L \mapsto {}^{\mathfrak{a}}L$ resp. $L \mapsto \overline{L}$ given above, these examples illustrate that there are cases in which there cannot exist any bijection preserving the algebraic structure, i.e. for example the decomposition type or the unitary groups. However, for ternary lattices we can explain the existence of some bijection.

Proposition 3.9. *For definite unimodular lattices of odd rank we have that either $h_{[\mathfrak{a}]} = h_{[\mathfrak{b}]}$ for all $[\mathfrak{a}], [\mathfrak{b}] \in \mathcal{C}_K$ or there is a violation to this inside of each ideal genus.*

Proof. Let $[\![\mathfrak{a}]\!], [\![\mathfrak{b}]\!]$ be distinct ideal genera. Then there exists $[\mathfrak{c}]$ such that $[\mathfrak{a}] = [\mathfrak{c}][\mathfrak{b}]$. In that case $[\![\mathfrak{c}^n\mathfrak{a}]\!] = [\![\mathfrak{c}\mathfrak{a}]\!] = [\![\mathfrak{b}]\!]$, since n is odd. Therefore multiplication by $[\mathfrak{c}]^n$ induces a bijection between $[\![\mathfrak{a}]\!]$ and $[\![\mathfrak{b}]\!]$. By Lemma 3.5 we therefore have a bijection of unimodular lattices with Steinitz class $[\mathfrak{a}'] \in [\![\mathfrak{a}]\!]$, and Steinitz class $[\mathfrak{b}'] \in [\![\mathfrak{b}]\!]$, so that $h_{[\mathfrak{a}']} = h_{[\mathfrak{b}']}$. So if $h_{[\mathfrak{a}']} = h_{[\mathfrak{a}]}$ holds for all $[\mathfrak{a}'] \in [\![\mathfrak{a}]\!]$, it holds for all classes in $\mathcal{C}_K$. On the converse if it is not satisfied for one ideal genus, it cannot be for any other. □

Corollary 3.10. *For definite unimodular lattices of rank 3 over K we have $h_{[\mathfrak{a}]} = h_{[\mathfrak{b}]}$ for all $[\mathfrak{a}], [\mathfrak{b}] \in \mathcal{C}_K$.*

Proof. From [HK89b] Theorem 5.2 we know that the special class numbers of all special classes contained in the genus $\mathbb{I}_{3,[\![\mathfrak{o}]\!]}$ are equal. By Corollary 1.3 we know that special and partial classes coincide in this case. Thus we have $h_{[\mathfrak{a}]} = h_{[\mathfrak{b}]}$ for all $[\mathfrak{a}], [\mathfrak{b}] \in [\![\mathfrak{o}]\!]$. Using Proposition 3.9 we get the claimed result. □

We have seen that for lattices of rank at most 3 we always have an equality of partial class numbers in a fixed genus (c.f. Corollaries 3.7 and 3.10). The underlying bijection quite often comes from some kind of algebraic construction (c.f. Lemmata 3.2 and 3.4), but for rank 3 not every such bijection is of that kind. In fact there are examples where there cannot exist any bijection preserving the order of the unitary groups or the decomposition types (c.f. Example 3.8). We have also seen that for rank 4 there is a counter example to the existence of any bijection at all (c.f. Example 3.1).
Due to the increasing complexity of the necessary computations to enumerate genera of rank ≥ 5, we have not found any more counter examples by the computational approach. Indeed, in every considered case the partial class numbers coincide. For example the genus $\mathbb{I}_{6,[\![\mathfrak{o}]\!]}$ over $\mathbb{Q}(\sqrt{-23})$ is a case not covered by the above results, where the partial class numbers of the three partial genera is 2965, but the sequences of orders of the unitary groups of the partial genera match only for two of the occuring Steinitz classes. Whether this phenomenon occurs only by chance or only for lattices of small rank, or if there is a more profound reason to be unveiled, remains an open question to us.

4. A mass formula for partial genera

Theorem 4.1. *Let $\mathcal{G}$ be a genus of definite unimodular lattices over K with $\mathcal{G}_{[\mathfrak{a}]} \subseteq \mathcal{G}$. Then the following holds:*

$$\operatorname{mass}(\mathcal{G}) = \frac{h_K}{2^{t-1}} \cdot \operatorname{mass}(\mathcal{G}_{[\mathfrak{a}]})$$

In particular, the partial masses are equal for all classes in the ideal genus $[\![\mathfrak{a}]\!]$.

Proof. Let $L_1, \ldots, L_{h_{[\mathfrak{a}]}}$ be a set of representatives for the classes in $\mathcal{G}_{[\mathfrak{a}]}$. By Lemma 1.1 the class of any L_i is the disjoint union of exactly $e(L_i) = [\mathfrak{o}^* : \det U(L_i)]$ special classes.
Now let $\Lambda_1 \ldots, \Lambda_m$ ($m = \sum_i e(L_i)$) be a set of representatives of the special classes contained in $\mathcal{G}_{[\mathfrak{a}]}$. By Lemma 1.2 $\mathcal{G}_{[\mathfrak{a}]}$ is the disjoint union of exactly $e(\mathcal{G})$ special genera. Now let $\mathcal{G}^+$ be the special genus of L_1 and assume that $\Lambda_1 \ldots, \Lambda_{h^+}$ is a set of representatives for the special classes in $\mathcal{G}^+$. Since $\det : U(L_i) \to \mathfrak{o}^*$ is a group homomorphism with kernel $SU(L_i)$, we get $|\det U(L_i)| = [U(L_i) : SU(L_i)]$. This gives

$$\frac{e(L_i)}{|SU(L_i)|} = \frac{|\mathfrak{o}^*|}{|\det U(L_i)| \cdot |SU(L_i)|} = \frac{|\mathfrak{o}^*|}{|U(L_i)|}.$$

Furthermore, since for isometric lattices $L_i \cong \Lambda_j$ the groups $SU(L_i)$ and $SU(\Lambda_j)$ are conjugate (in $U(V)$), the groups $SU(\Lambda_j)$ have the same order $|SU(L_i)|$ for all j with $L_i \cong \Lambda_j$. Thus we get

$$\begin{aligned} |\mathfrak{o}^*| \cdot \operatorname{mass}(\mathcal{G}_{[\mathfrak{a}]}) &= |\mathfrak{o}^*| \cdot \sum_{i=1}^{h_{[\mathfrak{a}]}} \tfrac{1}{|U(L_i)|} = \sum_{i=1}^{h_{[\mathfrak{a}]}} \tfrac{e(L_i)}{|SU(L_i)|} \\ &= \sum_{i=1}^{h_{[\mathfrak{a}]}} \sum_{\substack{j=1 \\ \Lambda_j \cong L_i}}^{m} \tfrac{1}{|SU(\Lambda_j)|} = \sum_{j=1}^{m} \tfrac{1}{|SU(\Lambda_j)|} \end{aligned} \tag{3}$$

Since $\Lambda_1, \ldots, \Lambda_m$ is a set of representatives of the special classes contained in $\mathcal{G}_{[\mathfrak{a}]}$, the last sum in (3) is the sum of the special masses of all special genera contained in $\mathcal{G}_{[\mathfrak{a}]}$. By [Reh71] (4.1) we get that the special mass does not depend on the special genus $\mathcal{G}^+$ but only on the genus $\mathcal{G}$. Since there are exactly $e(\mathcal{G})$ special genera contained in $\mathcal{G}_{[\mathfrak{a}]}$ all of them having the same special mass, we get the equation

$$\operatorname{mass}(\mathcal{G}_{[\mathfrak{a}]}) = \tfrac{e(\mathcal{G})}{|\mathfrak{o}^*|} \cdot \operatorname{mass}^+(\mathcal{G}^+).$$

Therefore also the partial mass does not depend on the Steinitz class $[\mathfrak{a}]$ and since there are exactly $\frac{h_K}{2^{t-1}}$ ideal classes in the ideal genus $[\![\mathfrak{a}]\!]$ we get the claimed formula. □

Using results of Gan and Yu (c.f. [GY00]) and Cho (c.f. [Cho12]) we now want to give an exact mass formula for the genera $\mathbb{I}_{n,[\![\mathfrak{a}]\!]}$ and $\mathbb{III}_{n,[\![\mathfrak{a}]\!]}$ described in chapter 2. The general mass formula for a genus $\mathcal{G}$ of definite unimodular lattices of rank n over K reads as follows (c.f. [GY00] Theorem 10.20):

$$\operatorname{mass}(\mathcal{G}) = 2 \cdot |d_K|^{\frac{n(n+1)}{4}} \cdot \prod_{j=1}^{n} \frac{(j-1)!}{(2\pi)^j} \cdot \prod_{p \in \mathbb{P}} \alpha_p(\mathcal{G})^{-1}, \tag{4}$$

where $\alpha_p(\mathcal{G})$ denotes the local density of the genus $\mathcal{G}$ at p. The local density of a unimodular lattice L for $p \in \mathbb{P}$ is given by Theorem 7.3. in [GY00] (we use the following convention: $\chi_K^i = 1 \Leftrightarrow i$ even and $\chi_K^i = \chi_K \Leftrightarrow i$ odd):

$$\alpha_p(\mathcal{G}) = \prod_{i=1}^{n} (1 - \chi_K^i(p) \cdot p^{-i}) \cdot \begin{cases} 1 & p \text{ unramified}, \\ 2 \cdot \lambda_p(\mathcal{G})^{-1} & p \text{ ramified}, \end{cases}$$

where for ramified primes $p \in \mathbb{P}$

$$\lambda_p(\mathcal{G}) := \begin{cases} 1 & n \text{ odd}, \\ 1 + (N(\mathfrak{a}), K/\mathbb{Q})_2 \cdot ((-1)^{\frac{p-1}{2}} p)^{-\frac{n}{2}} & n \text{ even}, p \neq 2, \\ 1 - 2^{-n} & n \text{ even}, p = 2, \mathcal{G} \text{ odd}, \\ 2^{1-n} & n \text{ even}, p = 2, \mathcal{G} \text{ even}, d_K \equiv_8 4, \\ 2^{-n}(1 + (N(\mathfrak{a}), K/\mathbb{Q})_2 \cdot ((-1)^{\frac{D-2}{4}} \cdot 2)^{-\frac{n}{2}}) & n \text{ even}, p = 2, \mathcal{G} \text{ even}, d_K \equiv_8 0. \end{cases} \tag{5}$$

In terms of Dirichlet L-series attached to χ_K^j the product of the inverse local densities $\alpha_p(\mathcal{G})^{-1}$ over all $p \in \mathbb{P}$ can be written as:

$$\prod_{p \in \mathbb{P}} \alpha_p(\mathcal{G})^{-1} = 2^{-t} \cdot \prod_{j=1}^{n} L(j, \chi_K^j) \cdot \prod_{p \mid d_K} \lambda_p(\mathcal{G}) \tag{6}$$

For further use, we state:

Proposition 4.2. *Let $\mathcal{G}$ be a genus of definite unimodular lattices of rank n over K with $\mathcal{G}_{[\mathfrak{a}]} \subseteq \mathcal{G}$. Then*

$$\operatorname{mass}(\mathcal{G}_{[\mathfrak{a}]}) = \frac{1}{|\mathfrak{o}^*|} \cdot |d_K|^{\frac{n^2+n-2}{4}} \cdot \prod_{j=2}^{n} \frac{(j-1)!}{(2\pi)^j} \cdot \prod_{j=2}^{n} L(j, \chi_K^j) \cdot \prod_{p \mid d_K} \lambda_p(\mathcal{G})$$

where $\lambda_p(\mathcal{G})$ is definied in (5).

Proof. Combining formulas (4) and (6) yields the following mass formula for the genus $\mathcal{G}$:

$$\operatorname{mass}(\mathcal{G}) = 2^{1-t} \cdot |d_K|^{\frac{n(n+1)}{4}} \cdot \prod_{j=1}^{n} \frac{(j-1)!}{(2\pi)^j} \cdot \prod_{j=1}^{n} L(j, \chi_K^j) \cdot \prod_{p \mid d_K} \lambda_p(\mathcal{G})$$

Using Theorem 4.1 and the class-number formula $h_K = \frac{|\mathfrak{o}^*|}{2\pi}\sqrt{|d_K|} \cdot L(1, \chi_K)$ (c.f. [Zag81] §8 Satz 5) we get exactly the claimed result. □

Corollary 4.3. *Let $\mathcal{G}$ be a genus of definite unimodular lattices of rank n over K with $\mathcal{G}_{[\mathfrak{a}]} \subseteq \mathcal{G}$. Then*

$$\operatorname{mass}(\mathcal{G}_{[\mathfrak{a}]}) = \frac{1}{|\mathfrak{o}^*|} \cdot \prod_{j=2}^{n} \frac{|B_{j,\chi_K^j}|}{2j} \cdot \prod_{p \mid d_K} \lambda_p(\mathcal{G}) \cdot \begin{cases} 1 & n \text{ odd,} \\ |d_K|^{n/2} & n \text{ even,} \end{cases}$$

where B_{j,χ_K^j} denote the genralized Bernoulli numbers and $\lambda_p(\mathcal{G})$ is definied in (5).

Proof. Using the functional equation for the L-series we get

$$|d_K|^{\frac{n(n+1)}{4}} \cdot \prod_{j=1}^{n} \frac{(j-1)!}{(2\pi)^j} \cdot \prod_{j=1}^{n} L(j, \chi_K^j) = 2^{-n} \cdot \prod_{j=1}^{n} L(1-j, \chi_K^j) \cdot \begin{cases} 1 & n \text{ odd,} \\ |d_K|^{n/2} & n \text{ even.} \end{cases}$$

Using $L(1-j, \chi_K^j) = \frac{|B_{j,\chi_K^j}|}{j}$ (c.f. [Was82]) and Theorem 4.1 we finally get the result. □

Remark 4.4. Proposition 4.3 makes the partial mass computable explicitly since the generalized Bernoulli numbers B_{j,χ_K^j} are given by the following formula (c.f. [Was82] Prop. 4.1)

$$B_{j,\chi_K^j} = \begin{cases} B_j(1) & j \text{ even,} \\ |d_K|^{j-1} \sum_{a=1}^{|d_K|} \chi_K(a) B_j(\frac{a}{|d_K|}) & j \text{ odd,} \end{cases}$$

where $B_j(X)$ denotes the usual Bernoulli polynomial.

5. Single-class partial genera

In this last section we determine all single-class partial genera of definite u9nimodular lattices of rank $n \geq 2$ over an imaginary-quadratic field. In particular we determine all single-class genera of such lattices. It turns out that a partial genus consists of a single isometry class if and only if the genus consists of a single isometry class.

Lemma 5.1. *Let $\mathcal{G}$ be a genus of definite unimodular lattices of rank n over K. The following inequality holds:*

$$\prod_{p \mid d_K} \lambda_p(\mathcal{G}) \geq 2^{-n} \cdot \frac{2^{n/2}-1}{(t+1)^{n/2}}.$$

Proof. It is easy to check by the definition of $\lambda_p(\mathcal{G})$ in (5) that:

$$\prod_{p|d_K} \lambda_p(\mathcal{G}) \geq 2^{-n} \cdot \prod_{p|d_K} \left(1 - p^{-\frac{n}{2}}\right) \tag{7}$$

If p_k denotes the kth prime number, we have $p_k \geq k+1$ and therefore

$$\prod_{p|d_K} \left(1 - p^{-\frac{n}{2}}\right) \geq \prod_{k=1}^{t} \left(1 - p_k^{-\frac{n}{2}}\right) \geq \prod_{k=1}^{t} \left(1 - (k+1)^{-\frac{n}{2}}\right) = \prod_{k=1}^{t} \frac{(k+1)^{\frac{n}{2}} - 1}{(k+1)^{\frac{n}{2}}}$$

The Bernoulli inequality gives us $\frac{(k+1)^{\frac{n}{2}}-1}{k^{\frac{n}{2}}} \geq 1$ for $k \in \{2, \ldots, t\}$ and therefore

$$\prod_{k=1}^{t} \frac{(k+1)^{\frac{n}{2}} - 1}{(k+1)^{\frac{n}{2}}} = (2^{\frac{n}{2}} - 1) \cdot \frac{3^{\frac{n}{2}} - 1}{2^{\frac{n}{2}}} \cdots \frac{(t+1)^{\frac{n}{2}} - 1}{t^{\frac{n}{2}}} \cdot \frac{1}{(t+1)^{\frac{n}{2}}} \geq \frac{2^{n/2} - 1}{(t+1)^{n/2}}$$

Together with (7) we get the claimed result. □

Proposition 5.2. *For all $n \geq 2$ there is a $d_{max}(n) \in \mathbb{N}$ such that for all single-class partial genera of definite unimodular lattices of rank n over K we have $|d_K| \leq d_{max}(n)$. For $2 \leq n \leq 7$ such a bound $d_{max}(n)$ is given by:*

n	2	3	4	5	6	7
$d_{max}(n)$	312	16	8	7	4	3

Proof. For a single-class partial genus $\mathcal{G}_{[\mathfrak{a}]}$ with representative L we have $\text{mass}(\mathcal{G}_{[\mathfrak{a}]}) = \frac{1}{|U(L)|}$. Since $|U(L)| \geq |\mathfrak{o}^*|$ for every lattice L, we necessarily have $\text{mass}(\mathcal{G}_{[\mathfrak{a}]}) \leq \frac{1}{|\mathfrak{o}^*|}$ for a single-class partial genus $\mathcal{G}_{[\mathfrak{a}]}$.
We now want to find a lower bound on the partial mass which for fixed rank $n \geq 2$ depends only on the discriminant d_K. Therefore we use the expression for the partial mass in Proposition 4.2. Since $\zeta_K(j) = \zeta(j) \cdot L(j, \chi_K)$ (c.f. [Zag81] p.100) and $\zeta_K(j) \geq 1$ for all $j > 1$ we easily deduce $L(j, \chi_K) = \frac{\zeta_K(j)}{\zeta(j)} \geq \frac{1}{\zeta(j)}$. Using $t \leq \log_2(|d_K|)$ and Lemma 5.1 for even n, it is easily seen that if

$$f_n(d) := \begin{cases} \frac{1}{|\mathfrak{o}^*|} \cdot d^{\frac{n^2+n-2}{4}} \cdot \prod_{j=2}^{n} \frac{(j-1)!}{(2\pi)^j} \cdot \frac{\zeta(2)\cdot\zeta(4)\cdot\ldots\cdot\zeta(n-1)}{\zeta(3)\cdot\zeta(5)\cdot\ldots\cdot\zeta(n)} & n \text{ odd}, \\ \frac{1}{|\mathfrak{o}^*|} \cdot d^{\frac{n^2+n-2}{4}} \cdot \frac{2^{n/2}-1}{2^{n(\log_2(|d_K|)+1)^{\frac{n}{2}}}} \cdot \prod_{j=2}^{n} \frac{(j-1)!}{(2\pi)^j} \cdot \frac{\zeta(2)\cdot\zeta(4)\cdot\ldots\cdot\zeta(n)}{\zeta(3)\cdot\zeta(5)\cdot\ldots\cdot\zeta(n-1)} & n \text{ even}, \end{cases}$$

we have $f_n(|d_K|) \leq \text{mass}(\mathcal{G}_{[\mathfrak{a}]})$ for all $n \in \mathbb{N}$. Since $f_n : \mathbb{N} \to \mathbb{R}_{\geq 0}$ is a strictly increasing function (in d) for $n \geq 2$, there is a $d_{\max}(n)$ such that $f_n(d) \leq \frac{1}{|\mathfrak{o}^*|}$ implies $d \leq d_{\max}(n)$. In particular we have $|d_K| \leq d_{\max}(n)$ if there is a single-class partial genus of definite unimodular lattices of rank n over K. For fixed n we can estimate this bound $d_{\max}(n)$. Using Corollary 4.3 we can compute the actual partial masses explicitly. Therefore, once given a bound $d_{\max}(n)$, we can try to improve it by substituting it by the greatest number $d \leq d_{\max}(n)$ such that d is a discriminant of some imaginary-quadratic field K and there is a partial genus of definite unimodular lattices of rank n over K with partial mass of the form $\frac{1}{2m}$ for some $m \in \mathbb{N}$. □

Instead of computing $d_{\max}(n)$ for all $n \geq 2$, we will effectively only need bounds for $n \in \{2, 4\}$. All other cases will be solved by constructing lattices with higher rank from other ones with lower rank. The procedure is as follows:

As seen in section 2, there is a free odd definite unimodular lattice $L^{\text{odd}}_{1,[\mathfrak{o}]}$ of rank 1 over every imaginary-quadratic field. Now the cancellation law implies that the mapping $L \mapsto L \perp L^{\text{odd}}_{1,[\mathfrak{o}]}$ is an injection preserving the Steinitz class. Thus, if the partial genus of odd definite unimodular lattices

of rank n consists of more than one single class, then the corresponding partial genus of odd definite unimodular lattices with the same Steinitz class and rank $n+1$ also consists of more than one class. In order to find all single-class partial genera we therefore have to determine the single-class partial genera of rank 2. Only for those we have to consider the corresponding partial genera of higher rank until we reach a partial genus consisting of more than one class.

We use essentially the same argument to determine all partial genera of even lattices. The procedure is slightly more difficult since even definite unimodular lattices exist only for even rank and there is a free even definite unimodular lattice $L^{\text{even}}_{4,[\mathfrak{o}]}$ of rank 4 for every imaginary-quadratic field (c.f. Corollary 2.4), but not necessarily of rank 2. Therefore we have to determine all single-class partial genera of even definite unimodular lattices rank 2 and 4 to proceed as above.

To check whether a partial genus consists of only one class, we compute the actual partial mass using Corollary 4.3. If this mass is of the form $\frac{1}{2m}$ for some $m \in \mathbb{N}$, the corresponding partial genus has to be tested further. For such a remaining partial genus $\mathcal{G}_{[\mathfrak{a}]}$ we construct a lattice L in it using the proof of Theorem 2.3. Then $\mathcal{G}_{[\mathfrak{a}]}$ is a single-class partial genus if and only if $\text{mass}(\mathcal{G}_{[\mathfrak{a}]}) = \frac{1}{|U(L)|}$. Doing this we get a complete list of all single-class partial genera of definite unimodular lattices of rank $n \geq 2$ over an imaginary-quadratic field. This proves the following Theorem:

Theorem 5.3. *For a genus $\mathcal{G}$ of definite unimodular lattices of rank $n \geq 2$ over an imaginary-quadratic field the following are equivalent:*

a) A partial genus $\mathcal{G}_{[\mathfrak{a}]}$ contained in $\mathcal{G}$ consists of only one isometry class.
b) $\mathcal{G}$ consists of only one isometry class.
c) $\mathcal{G}$ is included in the following list:

d_K	*genera*
-3	$\mathbb{I}_{2,[\![\mathfrak{o}]\!]}$, $\mathbb{I}_{3,[\![\mathfrak{o}]\!]}$, $\mathbb{I}_{4,[\![\mathfrak{o}]\!]}$, $\mathbb{I}_{5,[\![\mathfrak{o}]\!]}$
-4	$\mathbb{I}_{2,[\![\mathfrak{o}]\!]}$, $\mathbb{I}_{3,[\![\mathfrak{o}]\!]}$, $\mathbb{I}_{4,[\![\mathfrak{o}]\!]}$, $\mathbb{III}_{4,[\![\mathfrak{o}]\!]}$
-7	$\mathbb{I}_{2,[\![\mathfrak{o}]\!]}$
-8	$\mathbb{I}_{2,[\![\mathfrak{o}]\!]}$, $\mathbb{III}_{2,[\![\mathfrak{o}]\!]}$
-20	$\mathbb{I}_{2,[\![\mathfrak{a}]\!]}$
-24	$\mathbb{III}_{2,[\![\mathfrak{o}]\!]}$, $\mathbb{III}_{2,[\![\mathfrak{a}]\!]}$
-40	$\mathbb{III}_{2,[\![\mathfrak{o}]\!]}$, $\mathbb{III}_{2,[\![\mathfrak{a}]\!]}$
-88	$\mathbb{III}_{2,[\![\mathfrak{a}]\!]}$

In each case $[\mathfrak{a}]$ is the unique non-trivial ideal class, and $[\![\mathfrak{a}]\!]$ denotes the unique non-trivial ideal genus.

□

Remark 5.4. As already mentioned, ideals and lattices of rank 1 happen to be essentially the same objects. Therefore the problem of finding all single class partial genera of definite unimodular lattices of rank $n = 1$ is trivial.
However the more familiar problem of finding all genera consisting of a single isometry class of lattices is in fact the problem on *idoneal numbers*: Finding all fundamental discriminants, such that each ideal genus consists of a single ideal class, i.e. $h_K = 2^{t-1}$. This is a long standing open problem and a solution lies well beyond the scope of the techniques used here.

References

[BCP97] W. Bosma, J. Cannon, and C. Playoust, *The Magma algebra system. I. The user language*, Journal of Symbolic Computation **24(3)** (1997), 235–265.

[Cho12] S. Cho, *Group schemes and local densities of ramified hermitian lattices in residue characteristic 2*, arXiv:1210.7894 (2012).

[Eic52] M. Eichler, *Note zur Theorie der Kristallgitter*, Mathematische Annalen **152(1)** (1952), 51–55.

[GY00] W.T. Gan and J.K. Yu, *Group schemes and local densities*, Duke Mathematical Journal **105(3)** (2000), 497–524.

[HK89a] K. Hashimoto and H. Koseki, *Class numbers of definite unimodular hermitian forms over the rings of imaginary quadratic fields*, Tohoku Mathematical Journal **41(1)** (1989), 1–30.

[HK89b] ———, *Class numbers of positive definite binary and ternary unimodular hermitian forms*, Tohoku Mathematical Journal **41(2)** (1989), 171–216.

[Hof91] D.W. Hoffmann, *On positive definite hermitian forms*, manuscripta mathematica **71(1)** (1991), 399–429.

[Jac62] R. Jacobowitz, *Hermitian forms over local fields*, American Journal of Mathematics **84(3)** (1962), 441–465.

[O'M73] O.T. O'Meara, *Introduction to quadratic forms*, Springer, 1973.

[Reh71] U. Rehmann, *Klassenzahlen einiger totaldefiniter klassischer Gruppen über Zahlkörpern*, Ph.D. thesis, Göttingen, 1971.

[Sch85] W. Scharlau, *Quadratic and hermitian forms*, Springer, 1985.

[Sch98] A. Schiemann, *Classification of hermitian forms with the neighbour method*, Journal of Symbolic Computation **26(4)** (1998), 487–508.

[Was82] L.C. Washington, *Introduction to Cyclotomic Fields*, Springer, 1982.

[Zag81] D.B. Zagier, *Zetafunktionen und quadratische Körper: eine Einführung in die höhere Zahlentheorie*, Springer, 1981.

Fakultät für Mathematik, Technische Universität Dortmund, 44221 Dortmund, Germany
E-mail address: michael.juergens@math.tu-dortmund.de

Fakultät für Mathematik, Technische Universität Dortmund, 44221 Dortmund, Germany
E-mail address: marc-christian.zimmermann@math.tu-dortmund.de